

AD-A184 342

K-CO ON TRANSITION METALS: A LOCAL IONIC INTERACTION
(U) PENNSYLVANIA UNIV PHILADELPHIA DEPT OF PHYSICS
C H PATTERSON ET AL MAY 87 TR-5 N00014-86-K-0304


1/1

UNCLASSIFIED

F/G 7/2

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A184 342

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0304

R&T Code 413F0005

Technical Report No. 5

DTIC
ELECTE
SEP 01 1987
S D

K-CO on Transition Metals: A Local Ionic Interaction

by

Charles H. Patterson, Peter A. Schultz, and Richard P. Messmer

Abstract Submitted

for the Annual Meeting of the

Electrochemical Society

University of Pennsylvania
Department of Physics
Philadelphia, PA

May 1987

Reproduction in whole or in part is permitted for
any purpose of the United States Government

* This document has been approved for public release
and sale; its distribution is unlimited.

*This statement should also appear in Item 10 of the Document Control Data-DD
Form 1473. Copies of the form available from cognizant contract administrator.

87 9 1 011

K-CO on Transition Metals:
A Local Ionic Interaction

Charles H. Patterson and
Peter A. Schultz
Department of Physics
University of Pennsylvania
Philadelphia, PA 19104

Richard P. Messmer
General Electric
Corporate Research and
Development Center
Schenectady, NY 12301
and

Department of Physics
University of Pennsylvania
Philadelphia, PA 19104

We wish to address the nature of the K-CO interaction on a transition metal surface when the K:CO stoichiometry is ≤ 1 . The interaction proposed is transfer of the K 4s electron to CO. A Born-Haber cycle (Figure 1) for this process on a surface has been calculated. The reference state is K plus CO, bound separately to the surface (a). A key point in the cycle is that removal of the electron from K bound to a metal surface (b) is less costly in energy (metal/K surface, i.e. $\sim +2.0$ eV) than from K in the gas phase (IP = +4.3 eV). This means that the complex is significantly stabilized on the surface. The removed K electron is then transported to the isolated adsorbed CO (c) at a cost of the electron affinity of CO (+1.5 eV). When the isolated ions are brought together (d) the net stabilization at the equilibrium geometry is ~ -1.8 eV.

Generalized valence bond calculations have been performed for the KCO complex and we find that it resembles KCN in that it is predominantly ionic (A Mulliken population analysis indicates 97% electron transfer from K to CO) and has a similar K-CX bond length. The K-CO bond length was found to be 2.53 Å and the CO bond length was 1.27 Å (cf. 1.15 Å for CO in the gas phase). The CO stretching frequency was found to be 1525 cm^{-1} . Surface vibrational spectroscopies have usually found C-O stretching frequencies for K+CO coadsorption systems in the range 1300-1500 cm^{-1} . IR spectroscopy of matrix isolated KCO found a C-O stretching frequency of 1600 cm^{-1} (1) and EPR studies of the same system (1) showed that K had transferred an electron to CO in the complex. Evidence for the extreme weakening of the C-O bond by K on the surface is provided by thermal desorption: complete isotropic scrambling results when a mixture of $^{12}\text{C}^{16}\text{O}$ and $^{13}\text{C}^{16}\text{O}$ is adsorbed with K present (2). A local interaction is indicated by the fact that K and CO desorb in coincident peaks from the coadsorbate system which are not observed for single adsorbate experiments (3).




Figure 1 - Born-Haber cycle for formation of K^+CO^- complex.

(1) R.H. Hauge and J.L. Margrave, Argonne National Lab (Report) ANL 1977, ANL-77-2. Conf. High Temp. Sci. Open-cycle Coal Fired MHD Syst. pp. 283-9.

(2) R.A. dePaola, J. Hrbek and F.M. Hoffmann, J. Chem. Phys. **82**, 2484 (1985).

(3) R.A. dePaola, J. Hrbek and F.M. Hoffmann, Chem. Phys. Lett. **106**, 83 (1984).



A-1

END

10-87

DTIC